



Egyptian Mathematical Society
Journal of the Egyptian Mathematical Society

www.etms-eg.org
www.elsevier.com/locate/joems

**ORIGINAL ARTICLE**

Solutions of some class of nonlinear PDEs in mathematical physics



Shoukry El-Ganaini

Department of Mathematics, Faculty of Science, Damanshour University, Bahira 22514, Egypt

Received 7 January 2015; revised 14 February 2015; accepted 17 February 2015

Available online 8 May 2015

Keywords

Modified simple equation method;
 Traveling wave solution;
 Nonlinear partial differential equation

Abstract In this work, the modified simple equation (MSE) method is applied to some class of nonlinear PDEs, namely, a system of nonlinear PDEs, a $(2 + 1)$ -dimensional nonlinear model generated by the Jaulent–Miodek hierarchy, and a generalized KdV equation with two power nonlinearities.

As a result, exact traveling wave solutions involving parameters have been obtained for the considered nonlinear equations in a concise manner. When these parameters are chosen as special values, the solitary wave solutions are derived. It is shown that the proposed technique provides a more powerful mathematical tool for constructing exact solutions for a broad variety of nonlinear PDEs in mathematical physics.

Mathematics Subject Classification: 35C05; 35L05; 35Q99

Copyright 2015, Egyptian Mathematical Society. Production and hosting by Elsevier B.V.

This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

1. Introduction

Recently, several methods have been used to extract exact solutions of nonlinear partial differential equations (NPDEs) such as inverse scattering method [1,2], the Darboux transform [3,4], the Hirota bilinear method [5,6], the Backlund transformation method [7–9], the Exp-function method [10–12], the (G'/G) -expansion method [13–15], the projective Riccati equation method [16], first integral method [17–20], exp $(-\Phi(\xi))$ -expansion method [21–23], the functional variable

method [24,25], modified simple equation method [26–33] and others. However, up to now, a unified method that can be used to deal with all types of nonlinear PDEs has not been found yet. To obtain more different types of exact solution, the enhancement of these methods is a challenge topic.

This paper is organized as follows: a description of the modified simple equation (MSE) method is presented first [26–33]. This is followed by an application of this method to three distinct model equations, two of them are $(2 + 1)$ -dimensional nonlinear types, namely, a system of nonlinear PDEs and a nonlinear model generated by Jaulent–Miodek hierarchy, and the third is the generalized KdV nonlinear model with two power nonlinearities.

Finally, some conclusions are given.

Peer review under responsibility of Egyptian Mathematical Society.



Production and hosting by Elsevier

2. The MSE method

Consider a general nonlinear PDE in the form of

$$P(u, u_t, u_x, u_{xx}, u_{tt}, u_{xt}, u_{xxx}, \dots) = 0. \quad (1)$$

where P is a polynomial in its arguments.

Step 1. Seek solitary wave solutions of Eq. (1) by taking

$$u(x, t) = U(z), \quad z = x - ct + \zeta, \quad (2)$$

where ζ is an arbitrary constant, and transform (1) to a nonlinear ordinary differential equation (ODE) as

$$Q(U, U', U'', U''', \dots) = 0. \quad (3)$$

where the prime denotes the derivation with respect to z .

Step 2. We suppose that (3) has the formal solution as

$$U(z) = \sum_{i=0}^M \alpha_i \left(\frac{\Psi'(z)}{\Psi(z)} \right)^i, \quad (4)$$

where $\alpha_i, (i = 0, 1, \dots, M)$ are constants to be determined such that $\alpha_M \neq 0$. The function $\Psi(z)$ is an unknown function to be determined later such that $\Psi'(z) \neq 0$.

Step 3. We determine the positive integer M in (4) by balancing the highest order derivatives and the nonlinear terms in (3).

Step 4. We substitute (4) into (3), we calculate all the necessary derivatives U', U'', U''', \dots and then we account the function $\Psi(z)$.

As a result of this substitution, we get a polynomial of $\Psi^{-j}(z) (j = 0, 1, \dots)$, with the derivatives of $\Psi(z)$. Equating all the coefficients of $\Psi^{-j}(z) (j = 0, 1, \dots)$, to zero yields a system of equations which can be solved to obtain α_i and $\Psi(z)$. Finally, substituting the values of α_i and $\Psi(z)$ and its derivative $\Psi'(z)$ into (4) leads to exact solutions of (1).

3. Applications

Example 1. A system of nonlinear PDEs given by [34–36] is

$$iu_t + n(u_{xx} + \alpha_1 u_{yy}) + \beta_1 |u|^2 u + \gamma_1 uv = 0 \quad (5)$$

$$\alpha_2 v_{tt} + (v_{xx} - \beta_2 v_{yy}) + \gamma_2 (|u|^2)_{xx} = 0 \quad (6)$$

where $n, \alpha_i, \beta_i, \gamma_i (i = 1, 2)$ are real constants and $n \neq 0, \beta_1 \neq 0, \gamma_1 \neq 0, \gamma_2 \neq 0$. The important cases of (5) and (6) are shown in [36–38] that are the nonlinear Schrodinger equation [36], the Davey–Stewartson (DS) equations [37], and the generalized Zakharov (GZ) equations [38].

At first we are going to extract exact solutions for (5) and (6), for this purpose, using the transformation formula

$$\begin{aligned} u(x, y, t) &= e^{i\theta} \varphi(\xi), v(x, y, t) = \Gamma(\xi), \\ \theta &= px + qy + \varepsilon t, \xi = x + cy + dt + \xi_0 \end{aligned} \quad (7)$$

where $p, q, \varepsilon, c, \xi_0$ and d are real constants.

Substituting (7) into (5) and (6), we can know that $d = -2n(p + \alpha_1 qc)$, and φ, Γ satisfy the equations

$$\begin{aligned} -(\omega + p^2 n + n\alpha_1 q^2) \varphi(\xi) + (n + n\alpha_1 c^2) \varphi''(\xi) + \beta_1 \varphi^3(\xi) \\ + \gamma_1 \varphi(\xi) \Gamma(\xi) = 0, \end{aligned} \quad (8)$$

$$(\alpha_2 d^2 - \beta_2 c^2 + 1) \Gamma''(\xi) + \gamma_2 (\varphi^2(\xi))'' = 0. \quad (9)$$

Integrating (9) twice with respect to ξ and considering the constant of integration equal to zero, we find

$$v(x, y, t) = \Gamma(\xi) = -\frac{\gamma_2}{(\alpha_2 d^2 - \beta_2 c^2 + 1)} \varphi^2(\xi). \quad (10)$$

Substituting (10) into (8), gives

$$\varphi''(\xi) - \lambda \varphi(\xi) - \mu \varphi^3(\xi) = 0, \quad (11)$$

where

$$\lambda = \frac{\omega + p^2 n + n\alpha_1 q^2}{n + n\alpha_1 c^2}, \quad \mu = \frac{-\beta_1 (\alpha_2 d^2 - \beta_2 c^2 + 1) + \gamma_1 \gamma_2}{(n + n\alpha_1 c^2)(\alpha_2 d^2 - \beta_2 c^2 + 1)}. \quad (12)$$

Balancing $\varphi''(\xi)$ with $\varphi^3(\xi)$ yields $M = 1$. Consequently, we have the formal solution as

$$\varphi(\xi) = A_0 + A_1 \left(\frac{\Psi'(\xi)}{\Psi(\xi)} \right), \quad (13)$$

where A_0 and A_1 are constants to be determined with $A_1 \neq 0$. Also the function $\Psi(\xi)$ has to be determined where $\Psi'(\xi) \neq 0$.

It is easy to see that

$$\varphi'(\xi) = A_1 \left(\frac{\Psi''}{\Psi} - \frac{\Psi'^2}{\Psi^2} \right), \quad (14)$$

$$\varphi''(\xi) = A_1 \left(\frac{\Psi'''}{\Psi} - 3 \frac{\Psi' \Psi''}{\Psi^2} + 2 \frac{\Psi'^3}{\Psi^3} \right). \quad (15)$$

Substituting (13)–(15) into (11) and equating all the coefficients of $\Psi^0, \Psi^{-1}, \Psi^{-2}, \Psi^{-3}$, to zero, we respectively obtain

$$\Psi^0: -\lambda A_0 - \mu A_0^3 = 0, \quad (16)$$

$$\Psi^{-1}: A_1 \Psi''' - \lambda A_1 \Psi' - 3A_0^2 A_1 \mu \Psi' = 0, \quad (17)$$

$$\Psi^{-2}: -3A_1 \Psi' \Psi'' - 3A_0 A_1^2 \mu \Psi'^2 = 0, \quad (18)$$

$$\Psi^{-3}: 2A_1 \Psi'^3 - \mu A_1^3 \Psi'^3 = 0, \quad (19)$$

From (16) and (19), we deduce that

$$A_0 = 0, \quad A_0 = \pm \sqrt{\lambda/\mu}, \quad A_1 = \pm \sqrt{2/\mu} \quad (20)$$

From (17) and (18), we obtain

$$\Psi''' / \Psi'' = -\ell_1, \quad (21)$$

where

$$\ell_1 = (\lambda + 3A_0^2 \mu) / (A_0 A_1 \mu). \quad (22)$$

Integrating (21), we obtain

$$\Psi'' = C_1 e^{-\ell_1 \xi}, \quad (23)$$

where C_1 is a constant of integration.

And from (18) and (23), we obtain

$$\Psi' = -m_1 e^{-\ell_1 \xi} \quad (24)$$

where

$$m_1 = C_1 / (A_0 A_1 \mu). \quad (25)$$

Integrating (24) with respect to ξ , we get

$$\Psi(\xi) = C_2 + (m_1 / \ell_1) e^{-\ell_1 \xi}, \quad (26)$$

where C_2 is the constant of integration.

Substituting the values of Ψ and Ψ' into (13), we obtain

$$\varphi(\xi) = A_0 - A_1 \left((m_1 e^{-\ell_1 \xi}) / \left(C_2 + \frac{m_1}{\ell_1} e^{-\ell_1 \xi} \right) \right). \quad (27)$$

Case 1. When $A_0 = 0$, solution (27) collapses and hence this case is rejected.

Case 2. When $A_0 = \pm\sqrt{\lambda/\mu}$, $A_1 = \pm\sqrt{2/\mu}$, where λ and μ are given in (12), substituting the values of A_0 and A_1 into (27) and simplifying, we obtain the exact solution as follows:

$$\begin{aligned} \varphi(x, y, t) = & \pm \sqrt{\frac{(\alpha_2 d^2 - \beta_2 c^2 + 1)(\omega + p^2 n + n\alpha_1 q^2)}{-\beta_1(\alpha_2 d^2 - \beta_2 c^2 + 1) + \gamma_1 \gamma_2}} \\ & \mp \frac{(n + n\alpha_1 c^2) \sqrt{\alpha_2 d^2 - \beta_2 c^2 + 1} C_1}{\sqrt{(-\beta_1(\alpha_2 d^2 - \beta_2 c^2 + 1) + \gamma_1 \gamma_2)(\omega + p^2 n + n\alpha_1 q^2)}} \\ & \times \left(\frac{e^{-\ell_1(x+cy+dt+\xi_0)}}{C_2 + \frac{(n+n\alpha_1 c^2)C_1}{2(\omega+p^2 n+n\alpha_1 q^2)} e^{-\ell_1(x+cy+dt+\xi_0)}} \right). \end{aligned} \quad (28)$$

Since C_1 and C_2 are arbitrary constants, therefore, if we set $C_1 = [2(\omega + p^2 n + n\alpha_1 q^2)] / (n + n\alpha_1 c^2)$, we obtain the exact solution of the system. (5) and (6) and can be written as

$$\begin{aligned} u(x, y, t) = & \pm \sqrt{\frac{(\alpha_2 d^2 - \beta_2 c^2 + 1)(\omega + p^2 n + n\alpha_1 q^2)}{-\beta_1(\alpha_2 d^2 - \beta_2 c^2 + 1) + \gamma_1 \gamma_2}} \\ & \times \left[1 \mp 2 \frac{e^{-\ell_1(x+cy+dt+\xi_0)}}{C_2 + e^{-\ell_1(x+cy+dt+\xi_0)}} \right] \\ & \times \exp[i(px + qy + \varepsilon t)], \end{aligned} \quad (29)$$

$$\begin{aligned} v(x, y, t) = & \frac{\gamma_2(\omega + p^2 n + n\alpha_1 q^2)}{-\beta_1(\alpha_2 d^2 - \beta_2 c^2 + 1) + \gamma_1 \gamma_2} \\ & \times \left[1 \mp 2 \frac{e^{-\ell_1(x+cy+dt+\xi_0)}}{C_2 + e^{-\ell_1(x+cy+dt+\xi_0)}} \right]^2. \end{aligned} \quad (30)$$

Therefore, the exact solution (29) and (30) turns out to the following solitary wave solution:

$$\begin{aligned} u_1(x, y, t) = & \pm \sqrt{\frac{(\alpha_2 d^2 - \beta_2 c^2 + 1)(\omega + p^2 n + n\alpha_1 q^2)}{-\beta_1(\alpha_2 d^2 - \beta_2 c^2 + 1) + \gamma_1 \gamma_2}} \\ & \times \tanh\left(\frac{1}{2} \sqrt{\frac{\omega + p^2 n + n\alpha_1 q^2}{2(n + n\alpha_1 c^2)}} (x + cy + dt + \xi_0)\right) \\ & \times \exp[i(px + qy + \varepsilon t)], \end{aligned} \quad (31)$$

$$\begin{aligned} v_1(x, y, t) = & \frac{\gamma_2(\omega + p^2 n + n\alpha_1 q^2)}{-\beta_1(\alpha_2 d^2 - \beta_2 c^2 + 1) + \gamma_1 \gamma_2} \\ & \times \tanh^2\left(\frac{1}{2} \sqrt{\frac{\omega + p^2 n + n\alpha_1 q^2}{2(n + n\alpha_1 c^2)}} (x + cy + dt + \xi_0)\right), \end{aligned} \quad (32)$$

when $C_2 = 1$.

On the other hand, if we set $C_2 = -1$, solution (29) and (30) reduces to the solitary wave solution as follows:

$$\begin{aligned} u_2(x, y, t) = & \pm \sqrt{\frac{(\alpha_2 d^2 - \beta_2 c^2 + 1)(\omega + p^2 n + n\alpha_1 q^2)}{-\beta_1(\alpha_2 d^2 - \beta_2 c^2 + 1) + \gamma_1 \gamma_2}} \\ & \times \coth\left(\frac{1}{2} \sqrt{\frac{\omega + p^2 n + n\alpha_1 q^2}{2(n + n\alpha_1 c^2)}} (x + cy + dt + \xi_0)\right) \\ & \times \exp[i(px + qy + \varepsilon t)]. \end{aligned} \quad (33)$$

$$\begin{aligned} v_2(x, y, t) = & \frac{\gamma_2(\omega + p^2 n + n\alpha_1 q^2)}{-\beta_1(\alpha_2 d^2 - \beta_2 c^2 + 1) + \gamma_1 \gamma_2} \\ & \times \coth^2\left(\frac{1}{2} \sqrt{\frac{\omega + p^2 n + n\alpha_1 q^2}{2(n + n\alpha_1 c^2)}} (x + cy + dt + \xi_0)\right). \end{aligned} \quad (34)$$

Comparing these results with the results obtained in [34–36], it can be seen that the solutions here are new.

Example 2. A $(2 + 1)$ -dimensional nonlinear model generated by the Jaulent–Miodek hierarchy [39] is

$$u_t = -\frac{1}{4}(u_{xx} - 2u^3)_x - \frac{3}{4}\left(\frac{1}{4}\partial_x^{-1}u_{yy} + u_x\partial_x^{-1}u_y\right), \quad (35)$$

where ∂_x^{-1} is the inverse of ∂_x with $\partial_x\partial_x^{-1} = \partial_x^{-1}\partial_x = 1$ and $(\partial_x^{-1}f)(x) = \int_{-\infty}^x f(t)dt$, under the decaying condition at infinity.

This model was studied by Liu and Yan in [40] using qualitative analysis method. Wazwaz in [39] has derived the multiple kink solutions and multiple singular kink solutions for this model. In addition, in [41], Taha and Noorani have obtained the exact solutions of (35) by using the (G'/G) -expansion method.

Now, we conduct our analysis to this model to extract its exact solutions. For this purpose, we introduce the potential

$$u(x, y, t) = v_x(x, y, t), \quad (36)$$

that transforms (35) to

$$v_{xt} + \frac{1}{4}v_{xx}v_{xx} - \frac{3}{2}v_x^2v_{xx} + \frac{3}{16}v_{yy} + \frac{3}{4}v_{xx}v_y = 0 \quad (37)$$

Let $v(x, y, t) = v(\xi)$, $\xi = x + y + ct + \xi_0$, where c is the wave speed and ξ_0 is an arbitrary constant. Putting the wave variable $\xi = x + y + ct + \xi_0$ into (37) yields

$$c v'' + \frac{1}{4}v''' - \frac{3}{2}(v')^2 v'' + \frac{3}{16}v'' + \frac{3}{4}v''v' = 0. \quad (38)$$

Integrating (38) with respect to ξ once and considering the constant of integration to be zero, we find

$$16 c v' + 4v'' - 8(v')^3 + 3v' + 6(v')^2 = 0 \quad (39)$$

Assuming that

$$v' = w, \quad (40)$$

and substituting (40) into (39), we obtain

$$16 c w + 4w'' - 8w^3 + 3w + 6w^2 = 0 \quad (41)$$

As in the previous example, we assume the following solution formula:

$$w(\xi) = A_0 + A_1 \left(\frac{\Psi'(\xi)}{\Psi(\xi)} \right), \quad (42)$$

where A_0 and A_1 are constants to be determined and $A_1 \neq 0$. Thus we obtain the following system of equations

$$\Psi^0 : 16cA_0 - 8A_0^3 + 3A_0 + 6A_0^2 = 0, \quad (43)$$

$$\Psi^{-1} : 16cA_1\Psi' + 4A_1\Psi''' - 24A_0^2A_1\Psi' + 3A_1\Psi' + 12A_0A_1\Psi' = 0, \quad (44)$$

$$\Psi^{-2} : -12A_1\Psi'\Psi'' - 24A_0A_1^2\Psi'^2 + 6A_1^3\Psi'^2 = 0, \quad (45)$$

$$\Psi^{-3} : 8A_1\Psi'^3 - 8A_1^3\Psi'^3 = 0 \quad (46)$$

From (43) and (46), we find

$$A_0 = 0, \quad 8A_0^2 - 6A_0 - (16c + 3) = 0, \quad A_1 = \pm 1, \quad (47)$$

From (44) and (45), we get

$$\Psi'''/\Psi'' = -\ell_2, \quad (48)$$

where

$$\ell_2 = (16c - 24A_0^2 + 3 + 12A_0)/(2A_1^3 - 8A_0A_1^2). \quad (49)$$

Integrating (48), we obtain

$$\Psi'' = C_1 e^{-\ell_2 \xi}, \quad (50)$$

where C_1 is a constant of integration.

From (45) and (50), we obtain

$$\Psi' = m_2 e^{-\ell_2 \xi} \quad (51)$$

where

$$m_2 = 2C_1/(A_1^2 - 4A_1A_0). \quad (52)$$

Integrating (51) with respect to ξ , we get

$$\Psi(\xi) = C_2 - (m_2/\ell_2)e^{-\ell_2 \xi}, \quad (53)$$

where C_2 is the constant of integration.

Substituting the values of Ψ and Ψ' into (42), we obtain

$$w(\xi) = A_0 + A_1 \left((m_2 e^{-\ell_2 \xi}) / \left(C_2 - \frac{m_2}{\ell_2} e^{-\ell_2 \xi} \right) \right). \quad (54)$$

Case 1. When $A_0 = 0$, $A_1 \neq 0$, and $\Psi' \neq 0$, then the exact solution of (35) has the following formula:

$$w(\xi) = \left(\frac{3+16c}{2} \right) \left(\frac{e^{-\ell_2 \xi}}{C_2 - e^{-\ell_2 \xi}} \right). \quad (55)$$

If $C_2 = \pm 1$, the exact solutions (55) are transformed to the following formulas:

$$w_1(\xi) = \left(\frac{3+16c}{4} \right) \left(1 - \coth \left(\pm \frac{3+16c}{4} \xi \right) \right), \quad (56)$$

$$w_2(\xi) = - \left(\frac{3+16c}{4} \right) \left(1 - \tanh \left(\pm \frac{3+16c}{4} \xi \right) \right), \quad (57)$$

respectively.

But $w = dv/d\xi$, then $v = \int w d\xi$, and $u = \partial v / \partial x$, we find the exact solutions of (35) as

$$u_1(x, y, t) = \left(\frac{3+16c}{4} \right) \times \left[1 - \coth \left(\left(\frac{3+16c}{4} \right) (x + y + ct + \xi_0) \right) \right] \quad (58)$$

$$u_2(x, y, t) = \left(\frac{3+16c}{4} \right) \times \left[-1 \pm \tanh \left(\left(\frac{3+16c}{4} \right) (x + y + ct + \xi_0) \right) \right] \quad (59)$$

Case 2. When $A_0 \neq 0$, $A_1 \neq 0$, and $\Psi' \neq 0$, then we deduce from (44) and (42) that

$$(16c - 24A_0^2 + 3 + 12A_0)\Psi' + 4\Psi''' = 0 \quad (60)$$

$$(4A_0 - A_1)A_1\Psi' + 2\Psi'' = 0 \quad (61)$$

With the help of (47), we can simplify (60) to take the form

$$(3 - 8A_0)A_0\Psi' + 2\Psi'' = 0. \quad (62)$$

From (61) and (62), we have

$$\Psi'''/\Psi'' = \ell_3 \quad (63)$$

where $\ell_3 = (8A_0^2 - 3A_0)/[A_1(A_1 - 4A_0)]$. Integrating (63) and using (61), we deduce that

$$\Psi' = m_3 e^{\ell_3 \xi} \quad (64)$$

where $m_3 = 2C_1/[A_1(A_1 - 4A_0)]$, and then

$$\Psi = C_2 + (m_3/\ell_3)e^{\ell_3 \xi}, \quad (65)$$

where C_1 and C_2 are constants of integration. From (42), (64) and (65), we have the exact solution

$$w(\xi) = A_0 \left[1 + \left(\frac{8A_0 - 3}{A_1 - 4A_0} \right) \frac{e^{\ell_3 \xi}}{C_2 + e^{\ell_3 \xi}} \right], \quad (66)$$

$$\text{where } A_0 = \frac{3 \pm \sqrt{9+8(3+16c)}}{8}.$$

If $C_2 = \pm 1$, the solution (66) turns out to the following formulas:

$$w_3(\xi) = A_0 \left[1 + \left(\frac{8A_0 - 3}{2A_1 - 8A_0} \right) \left\{ 1 + \tanh \left(\frac{A_0(8A_0 - 3)}{(2A_1^2 - 8A_0A_1)} \xi \right) \right\} \right] \quad (67)$$

$$w_4(\xi) = A_0 \left[1 + \left(\frac{8A_0 - 3}{2A_1 - 8A_0} \right) \left\{ 1 + \coth \left(\frac{A_0(8A_0 - 3)}{(2A_1^2 - 8A_0A_1)} \xi \right) \right\} \right] \quad (68)$$

But $w = dv/d\xi$, then $v = \int w d\xi$, and $u = \partial v / \partial x$, we then find the exact solutions of (35) and can be written as

$$u_3(x, y, t) = A_0 + A_0 \left(\frac{8A_0 - 3}{\pm 2 - 8A_0} \right) \times \left\{ 1 + \tanh \left(\frac{A_0(8A_0 - 3)}{(2 \mp 8A_0)} (x + y + ct + \xi_0) \right) \right\} \quad (69)$$

$$u_4(x, y, t) = A_0 + A_0 \left(\frac{8A_0 - 3}{\pm 2 - 8A_0} \right) \times \left\{ 1 + \coth \left(\frac{A_0(8A_0 - 3)}{(2 \mp 8A_0)} (x + y + ct + \xi_0) \right) \right\} \quad (70)$$

respectively, where $A_0 = \frac{3 \pm \sqrt{33+128c}}{8}$, as given in (47).

Eqs. (58), (59) and (69), (70), are new types of exact traveling wave solutions to the (2 + 1)-dimensional nonlinear model generated by the Jaulent–Miodek hierarchy (35). It could not be obtained by the methods presented in [39–41].

Example 3. A generalized KdV equation with two power nonlinearities [42] is

$$u_t + (au^n - bu^{2n})u_x + u_{xxx} = 0 \quad (71)$$

This equation describes the propagation of nonlinear long acoustic-type waves.

If the amplitude is not supposed to be small, (71) serves as an approximate model for the description of weak dispersive effects on the propagation of nonlinear waves along a characteristic direction [42,43].

The wave variable $\xi = x - ct + \xi_0$, and integrating once, (71) is converted to the ODE

$$-cu + \frac{a}{n+1}u^{n+1} - \frac{b}{2n+1}u^{2n+1} + u'' = 0, \quad (72)$$

Balancing u^{2n+1} with u'' in (72), we find that $M = 1/n$.

We use the following transformation:

$$u = v^{1/n} \quad (73)$$

and using (73) into (72) gives

$$-cn^2(2n+1)(n+1)v^2 + an^2(2n+1)v^3 - bn^2(n+1)v^4 + n(2n+1)(n+1)vv'' + (1-n^2)(2n+1)v^2 = 0. \quad (74)$$

Balancing vv'' with v^4 gives $M = 1$. The modified simple equation method admits the use of the finite expansion

$$v(\xi) = A_0 + A_1 \left(\frac{\Psi'(\xi)}{\Psi(\xi)} \right), \quad (75)$$

where A_0 and A_1 are constants to be determined and $A_1 \neq 0$.

With similar procedures made in previous examples, we have

$$\Psi^0: -cn^2(2n+1)(n+1)A_0 + an^2(2n+1)A_0^3 - bn^2(n+1)A_0^4 = 0 \quad (76)$$

$$\Psi^{-1}: -2A_0A_1cn^2(2n+1)(n+1)\Psi' + 3A_0^2A_1an^2(2n+1)\Psi' - 4A_0^3A_1bn^2(n+1)\Psi' + nA_0A_1(2n+1)(n+1)\Psi''' = 0 \quad (77)$$

$$\begin{aligned} \Psi^{-2}: & -cn^2(2n+1)(n+1)A_1^2\Psi'^2 + 3A_0A_1^2an^2(2n+1), \\ & \Psi'^2 - 6A_0^2A_1^2bn^2(n+1), \\ & \Psi'^2 - 3A_0A_1n(2n+1)(n+1)\Psi'\Psi'' + n(2n+1)(n+1)A_1^2\Psi'\Psi''' \\ & + (1-n^2)(2n+1)A_1^3\Psi'^2 = 0 \end{aligned} \quad (78)$$

$$\begin{aligned} \Psi^{-3}: & an^2(2n+1)A_1^3\Psi'^3 - 4A_0A_1^3bn^2(n+1)\Psi'^3 \\ & + 2A_0A_1n(2n+1)(n+1)\Psi'^3 - 3A_1^2n(2n+1)(n+1)\Psi'^2\Psi'' \\ & - 2A_1^2(1-n^2)(2n+1)\Psi''\Psi'^2 = 0 \end{aligned} \quad (79)$$

$$\begin{aligned} \Psi^{-4}: & -bn^2(n+1)A_1^4\Psi'^4 + 2A_1^2n(2n+1)(n+1)\Psi'^4 \\ & + A_1^2(1-n^2)(2n+1)\Psi'^4 = 0 \end{aligned} \quad (80)$$

Thus from (76) and (80), we get

$$\begin{aligned} A_0 &= 0, bn^2(n+1)A_0^2 - an^2(2n+1)A_0 \\ &+ cn^2(2n+1)(n+1) = 0, \\ A_1 &= \pm \frac{1}{n} \sqrt{\frac{2n^2+3n+1}{b}} \end{aligned} \quad (81)$$

Let us consider here the case $A_0 = 0$, $A_1 = \pm \frac{1}{n} \sqrt{\frac{2n^2+3n+1}{b}}$, and only left the case of $A_0 \neq 0$, for the reader.

Therefore from (77), we deduce that

$$2cn\Psi' - \Psi''' = 0 \quad (82)$$

The general solution of this equation is given by

$$\Psi(\xi) = C_0 + C_1 e^{r_1 \xi} + C_2 e^{r_2 \xi}, \quad (83)$$

where $r_{1,2} = \pm \sqrt{2cn}$ and $C_i (i = 0, 1, 2)$, are arbitrary constants.

Consequently, the exact solutions of (71) have the form of

$$u(x, t) = \left[\pm \frac{1}{n} \sqrt{\frac{2n^2+3n+1}{b}} \times \left(\frac{C_1 r_1 e^{r_1(x-ct+\xi_0)} + C_2 r_2 e^{r_2(x-ct+\xi_0)}}{C_0 + C_1 e^{r_1(x-ct+\xi_0)} + C_2 e^{r_2(x-ct+\xi_0)}} \right) \right]^{\frac{1}{n}} \quad (84)$$

Comparing the solutions (84) with those obtained in [42], we see that our solutions are new.

Remark. For $n = 1$, $a = 6$, $b = \pm 6$, the solutions (84) are the solutions for the two models for the Gardner equation:

$$u_t + 6uu_x \pm 6u^2u_x + u_{xxx} = 0,$$

that describes internal solitary waves in shallow seas. These models are the positive and negative Gardner equation depending on the sign of the cubic nonlinear term.

4. Conclusion

In this paper, we have successfully implemented and applied the modified simple equation (MSE) method to find the exact and solitary wave solutions for some class of nonlinear PDEs, via a system of nonlinear PDEs, a (2 + 1)-dimensional nonlinear model generated by the Jaulent–Miodek hierarchy, and a generalized KdV equation with two power nonlinearities. The method offers solutions with free parameters that may be important for explaining some physical phenomena in mathematical physics. The solitary wave solutions are obtained by setting appropriate values to the free parameters. The MSE method has been achieved without using any symbolic computations, since the method is simple, effective and can be applied to many other nonlinear PDEs, especially for equations with distinct power law nonlinearities that can be done in a future work.

Acknowledgment

The author would like to thank the referees for their useful comments that led to some improvements of the current paper.

References

- [1] M.J. Ablowitz, P.A. Clarkson, *Solitons, Nonlinear Evolution Equations and Inverse Scattering*, Cambridge University Press, Cambridge, 1991.

- [2] R. Beals, R.R. Coifman, Scattering and inverse scattering for 1st order system, *Commun. Pure. Appl. Math.* 37 (1984) 39–90.
- [3] V.B. Matveev, M.A. Salle, *Darboux Transformation and Solitons*, Springer, Berlin, 1991.
- [4] E.G. Fan, A family of completely integrable multi-Hamiltonian systems explicitly related to some celebrated equations, *J. Math. Phys.* 42 (2001) 4327–4344.
- [5] R. Hirota, J. Satsuma, Soliton solution of a coupled KdV equation, *Phys. Lett. A* 85 (1981) 407–408.
- [6] R. Hirota, Exact solution of the Korteweg–de Vries equation for multiple collisions of solitons, *Phys. Rev. Lett.* 27 (1971) 1192–1194.
- [7] R.M. Miura, *Backlund Transformation*, Springer, Berlin, 1978.
- [8] N.A. Kudryashov, Special polynomials associated with some hierarchies, *Phys. Lett. A* 372 (2008) 945–956.
- [9] C. Rogers, W.F. Shadwick, *Backlund Transformation*, Academic Press, New York, 1982.
- [10] A.M. Wazwaz, Generalized solitonary and periodic solutions for nonlinear partial differential equations by the Exp-function method, *Nonlinear Dyn.* 52 (2008) 1–9.
- [11] J.H. He, X.H. Wu, Exp-function method for nonlinear wave equations, *Chaos Solitons Fract.* 30 (2006) 700–708.
- [12] A. Bekir, A. Boz, Exact solutions for nonlinear evolution equations using Exp-function method, *Phys. Lett. A* 372 (2008) 1619–1625.
- [13] M. Mirzazadeh, M. Eslami, A. Biswas, Soliton solutions of the generalized Klein–Gordon equation by using (G'/G) -expansion method, *Comput. Appl. Math.* 33 (3) (2014) 831–839.
- [14] S. Zhang, J. Tong, W. Wang, A generalized (G'/G) -expansion method for the mKdV equation with variable coefficients, *Phys. Lett. A* 372 (2008) 2254–2257.
- [15] J. Zhang, X. Wei, Y. Lu, A generalized (G'/G) -expansion method and its applications, *Phys. Lett. A* 372 (2008) 3653–3658.
- [16] Z.Y. Yan, Generalized method and its application in the higher-order nonlinear Schrödinger equation in nonlinear optical fibres, *Chaos Solitons Fract.* 16 (2003) 759–766.
- [17] Shoukry Ibrahim Atia El-Ganaini, Travelling wave solutions to the generalized Pochhammer–Chree (PC) equations using the first integral method, *Math. Prob Eng.* <http://dx.doi.org/10.1155/2011/629760>.
- [18] Shoukry Ibrahim Atia El-Ganaini, New exact solutions of some nonlinear systems of partial differential equations using the first integral method, *Abst. Appl. Anal.* <http://dx.doi.org/10.1155/2013/693076>.
- [19] Shoukry Ibrahim Atia El-Ganaini, The first integral method to the nonlinear schrodinger equations in higher dimensions, *Abstract Appl. Anal.* <http://dx.doi.org/10.1155/2013/349173>.
- [20] M. Mirzazadeh, M. Eslami, Exact solutions of the Kudryashov–Sinelschikov equation and nonlinear telegraph equation via the first integral method, *Nonlin. Anal., Model. Control* 17 (4) (2012) 481–488.
- [21] Md. Sabur Uddin, Md. Nur Alam, S.M. Shamim Hossain, Md. Samiul Hasan, M. Ali Akbar, Some new exact travelling wave solutions to the $(3 + 1)$ -dimensional Zakharov–Kuznetsov equation and the Burgers equations via Exp $(-\Phi(\xi))$ – expansion method, *FMIA* 1 (1) (2014) 1–8.
- [22] K. Khan, M.A. Akbar, Application of exp $(-\Phi(\xi))$ – expansion method to find the exact solutions of modified Benjamin–Bona–Mahoney equation, *World Appl. Sci. J.* 24 (2013) 1373–1377.
- [23] K. Khan, M. Ali. Akbar, Exact traveling wave solutions of Kadomtsev–Petviashvili equation, *JEMS* (2014), doi:10.1016/j.joems.2014.03.010.
- [24] A. Nazarzadeh, M. Eslami, M. Mirzazadeh, Exact solutions of some nonlinear partial differential equations using functional variable method, *Pramana* 81 (2) (2013) 225–236.
- [25] M. Eslami, M. Mirzazadeh, Functional variable method to study nonlinear evolution equations, *Cent. Eur. J. Eng.* 3 (3) (2013) 451–458.
- [26] A.J.M. Jawad, M.D. Petkovic, A. Biswas, Modified simple equation, method for nonlinear evolution equations, *Appl. Math. Comput.* 217 (2010) 869–877.
- [27] K. Khan, M.A. Akbar, N. HJ, M. Ali, The modified simple equation method for exact and solitaty wave solutions of nonlinear evolution equation: the GZK–BBM equation and right-handed non-commutative Burgers equations, *ISRN Math. Phys.* <http://dx.doi.org/10.1155/2013/146704>.
- [28] E.M.E. Zayed, S.A. Hoda Ibrahim, Exact solutions of nonlinear evolution equations in mathematical physics using the modified simple equation method, *Chin. Phys. Lett.* 29 (2012) 060201–060204.
- [29] E.M.E. Zayed, A.H. Arnous, The modified simple equation method and its applications to $(2+1)$ -dimensional systems of nonlinear evolution equation, *Sci. Res. Essays* 8 (40) (2013) 1973–1982.
- [30] M. Akbari, The modified simplest equation method and its application, *Comput. Meth. Diff. Eq.* 1 (1) (2013) 71–77.
- [31] K. Khan, M.A. Akbar, M.N. Alam, Traveling wave solutions of the nonlinear Drinfel, d – Sokolov–Wilson equation and modified Benjamin–Bona–Mahoney equations, *J. Energy Math. Soc.* 21 (2013) 233–240.
- [32] N. Taghizadeh, M. Mirzazadeh, A. Samiei Paghaleh, J. Vahidi, Exact solutions of nonlinear evolution equations by using the modified simple equation method, *Ain Shams Eng. J.* 3 (2012) 321–325.
- [33] M. Eslami, M. Mirzazadeh, A. Biswas, Soliton solutions of the resonant nonlinear Schrödinger equation in optical fibres with time-dependent coefficients by simplest equation approach, *J. Modern Optics* 60 (19) (2013) 1627–1636.
- [34] Y. Zhou, M. Wang, T. Miao, The periodic wave solutions and solitary For a class of nonlinear partial differential equations, *Phys. Lett. A* 323 (2004) 77–88.
- [35] Y. Feng, D. Wang, et al., More solutions of the auxiliary equation to get the solutions for a class of nonlinear partial differential equations, *Commun. Nonlinear Sci. Numer. Simul.* 15 (2010) 3332–3338.
- [36] B. Lu, Coupling Backlund transformation of Riccati equation and division theorem method for traveling wave solutions of some class of NLPDEs, *Commun. Nonlinear Sci. Numer. Simul.* 17 (2012) 4626–4633.
- [37] A. Davey, K. Stewartson, On three dimensional packets of surface waves, *Proc. R. Soc. Lond. A* 338 (1974) 101–110.
- [38] B. Malomed, D. Anderson, et al., Dynamics of solitary waves in the Zakharov model equations, *Phys. Rev. E* 55 (1997) 962–968.
- [39] A.-M. Wazwaz, Multiple kink solutions and multiple singular kink solutions for $(2 + 1)$ -dimensional nonlinear models generated by the Jaulent–Miodek hierarchy, *Phys. Lett. A* 373 (2009) 1844–1846.
- [40] H. Liu, F. Yan, The bifurcation and exact traveling wave solutions for $(2 + 1)$ -dimensional nonlinear models generated by the Jaulent–Miodek hierarchy, *Int. J. Nonlinear Sci.* 11 (2) (2011) 200–205.
- [41] W.M. Taha, M.S.M. Noorani, Exact solutions of equation generated by the Jaulent –Miodek hierarchy by (G'/G) -expansion method, *Math. Prob Eng.* <http://dx.doi.org/10.1155/2013/392830>.
- [42] A.-M. Wazwaz, *Partial Differential Equations and Solitary Waves Theory*, Higher Education Press, Beijing, 2009 (Springer –Verlag Berlin Heidelberg).
- [43] A.V. Slyunyaev, EN. Pelinovski, Dynamics of large-amplitude solitons, *J. Exp. Theor. Phys.* 89 (1) (1999) 173–181.